




F/EP 03/03343 'a. U6. 201

10/509519 

INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

REC'D 27 JUN 2003

WIPO

PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.


In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c.; plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed



Dated

6 May 2003

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

Patents Act 1977
Section 16



0207450.8 E707581-1 G69803
F01/7700 0.00-0207450.8

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office
Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference

AWGP/JW/PG4792

2. Patent:

0207450.8

(The Pate.

28 MAR 2002

3. Full name, address and postcode of the or of each applicant (*underline all surnames*)

Glaxo Group Limited
Glaxo Wellcome House, Berkeley Avenue,
Greenford, Middlesex UB6 0NN, Great Britain

Patents ADP number (*if you know it*)

If the applicant is a corporate body, give the country/state of its incorporation

United Kingdom

473587003

4. Title of the invention

Novel Process

5. Name of your agent (*if you have one*)

Corporate Intellectual Property

"Address for service" in the United Kingdom to which all correspondence should be sent

(*including the postcode*)

GlaxoSmithKline
Corporate Intellectual Property CN925.1
980 Great West Road
BRENTFORD
Middlesex TW8 9GS

Patents ADP number (*if you know it*)

796982003

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or each of these earlier applications and (*if you know it*) the or each application number

Country Priority application number Date of filing
(*if you know it*) (*day / month / year*)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(*day / month / year*)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (*Answer yes if:*

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is named as an applicant, or
 - c) any named applicant is a corporate body
- See note (d)

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form
Description
Claim(s)
Abstract
Drawings

22

10. If you are also filing any of the following, state how many against each item.

Priority Documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

We request the grant of a patent on the basis of this application

Signature

H B Dawson

Date 28-Mar-02

12. Name and daytime telephone number of person to contact in the United Kingdom

H B Dawson 01279 644689

Warning

After an application for a Patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission unless an application has been filed at least six weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.


Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- For details of the fee and ways to pay please contact the Patent Office.

Novel Process

This invention relates to novel processes, in particular to processes for preparing certain morpholine derivatives.

- 5 Co-pending International Patent Application number PCT/GB01/04530 (Glaxo Group Limited) relates to certain morpholine urea derivatives of formula (I)



10

wherein:

- R^1 represents C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl- Y^1 -, heteroaryl- Y^1 -, aryl-(O)_t-aryl- Y^1 -, aryl-(O)_t-heteroaryl- Y^1 -, heteroaryl-(O)_t-aryl- Y^1 -, heteroaryl-(O)_t-heteroaryl- Y^1 -, aryl-SO₂- Y^1 -, C_{1-6} alkyl-G- Y^1 -, heteroaryl-G-aryl- Y^1 -, heteroaryl-G-heteroaryl- Y^1 -, $R^{17}O(CO)-C_{2-6}$ alkenyl- Y^1 -, $R^{17}NHCO- Y^1 -, $R^{17}NHCO- Y^1 -, C_{2-6} alkenyl- Y^1 -, C_{2-6} alkynyl- Y^1 -, aryl-O- Y^1 -, heteroaryl-O- Y^1 -, C_{1-6} alkyl-SO₂- Y^1 -, M- Y^1 -, J¹- Y^1 -, J¹-CO- Y^1 -, aryl-CO- Y^1 - or C_{3-8} cycloalkyl- Y^1 - or C_{3-8} cycloalkenyl- Y^1 -, which C_{2-6} alkynyl and C_{2-6} alkynyl- Y^1 may be optionally substituted with a -OR¹⁷ group, which C_{2-6} alkenyl may be optionally substituted by one or more -COOR¹⁷ groups and which cycloalkyl or cycloalkenyl may be optionally substituted by one or more hydroxyl or C_{1-6} alkyl groups;$$
- R^2 represents hydrogen or C_{1-6} alkyl optionally substituted by a hydroxy group;
- R^3 represents hydrogen or C_{1-6} alkyl;
- 25 or R^1 and R^2 may together with the nitrogen atom to which they are attached form a group of formula J² wherein said nitrogen atom substitutes for either X¹ or X²;
- t represents 0 or 1;
- X represents ethylene or a group of formula CR^eR^f wherein R^e and R^f
- 30 independently represent hydrogen or C_{1-4} alkyl or R^e and R^f may together with the carbon atom to which they are attached form a C_{3-8} cycloalkyl group;
- R^4 and R^5 independently represent hydrogen or C_{1-4} alkyl;
- Z represents a bond, CO, SO₂, CR¹⁰R⁷(CH₂)_n, (CH₂)_nCR¹⁰R⁷, CHR⁷(CH₂)_nO, CHR⁷(CH₂)_nS, CHR⁷(CH₂)_nOCO, CHR⁷(CH₂)_nCO, COCHR⁷(CH₂)_n
- 35 or SO₂CHR⁷(CH₂)_n;

R^6 represents C_{1-6} alkyl, C_{2-6} alkenyl, aryl, heteroaryl, aryl- C_{2-6} alkenyl-, -CN or a group of formula $-Y^2-J^3$;

R^7 represents hydrogen, C_{1-4} alkyl, $CONR^8R^9$ or $COOC_{1-6}$ alkyl;

a and b represent 1 or 2, such that $a+b$ represents 2 or 3;


5 G represents $-SO_2-$, $-SO_2NR^{18}-$, $-NR^{18}SO_2-$, $-NR^{18}CO-$, CO or $-CONR^{18}-$;

n represents an integer from 0 to 4;

M represents a C_{3-8} cycloalkyl or C_{3-8} cycloalkenyl group fused to a monocyclic aryl or monocyclic heteroaryl group;

J^1 , J^2 and J^3 independently represent a moiety of formula (K):

10



wherein X^1 represents oxygen, NR^{11} or sulphur, X^2 represents CH_2 , oxygen, NR^{12} or sulphur, m^1 represents an integer from 1 to 3 and m^2 represents an

15 integer from 1 to 3, provided that m^1+m^2 is in the range from 3 to 5, also provided that when both X^1 and X^2 represent oxygen, NR^{11} , NR^{12} or sulphur, m^1 and m^2 must both not equal less than 1. The moiety K is optionally substituted by one or more (eg. 1 or 2) $-Y^3$ -aryl, $-Y^3$ -heteroaryl, $-Y^3$ -CO-aryl, $-COC_{3-8}$ cycloalkyl, $-Y^3$ -CO-heteroaryl, $-C_{1-6}$ alkyl, $-Y^3$ - $COOC_{1-6}$ alkyl, $-Y^3$ - COC_{1-6} alkyl, $-Y^3$ -W, $-Y^3$ -CO-W, $-Y^3$ - $NR^{15}R^{16}$, $-Y^3$ - $CONR^{15}R^{16}$, hydroxy, oxo, $-Y^3$ - $SO_2NR^{15}R^{16}$, $-Y^3$ - SO_2C_{1-6} alkyl, $-Y^3$ - SO_2 aryl, $-Y^3$ - SO_2 heteroaryl, $-Y^3$ - $NR^{13}C_{1-6}$ alkyl, $-Y^3$ - $NR^{13}SO_2C_{1-6}$ alkyl, $-Y^3$ - $NR^{13}CONR^{15}R^{16}$, $-Y^3$ - $NR^{13}COOR^{14}$ or $-Y^3$ - $OCQNR^{15}R^{16}$ groups, and is optionally fused to a monocyclic aryl or heteroaryl ring;

20 R^8 , R^9 , R^{10} , R^{11} , R^{12} , R^{13} and R^{14} independently represent hydrogen or C_{1-6} alkyl; R^{15} and R^{16} independently represent hydrogen or C_{1-6} alkyl or R^{15} and R^{16} together with the nitrogen atom to which they are attached may form a morpholine, piperidine or pyrrolidine ring;

R^{17} and R^{18} independently represent hydrogen or C_{1-6} alkyl;


30 W represents a saturated or unsaturated, non-aromatic 5-7 membered ring containing between 1 and 3 heteroatoms selected from nitrogen, oxygen or sulphur, optionally substituted with one or more C_{1-6} alkyl, halogen or hydroxy groups;

Y^1 , Y^2 and Y^3 independently represent a bond or a group of formula $-(CH_2)_pCR^cR^d(CH_2)_q-$ wherein R^c and R^d independently represent hydrogen or C_{1-4} alkyl or R^c and R^d may together with the carbon atom to which they are attached form a C_{3-8} cycloalkyl group, and p and q independently represent an integer from 0 to 5 wherein $p+q$ is an integer from 0 to 5;

and salts and solvates thereof;

with the provisos that;

the compound of formula (I) is not a compound of formula (I)^a:




5

wherein R^{2'} represents hydrogen or lower alkyl (specifically C₁₋₄ alkyl); R^{3'} represents hydrogen; X' represents methylene or ethylene; a' and b' both represent 1; R^{4'} and R^{5'} both represent hydrogen; and wherein the moiety -Z'-R^{6'}


10 represents halobenzyl, and;

the compound of formula (I) is not a compound of formula (I)^b:




- 15 wherein R^{1''} represents a hydrogen atom, a C₁₋₆ alkyl group, a C₃₋₆ cycloalkyl group, a C₃₋₆ cycloalkylC₁₋₄ alkyl group, an aryl group or an arylC₁₋₄ alkyl group (particularly wherein aryl represents phenyl or naphthyl) in which the aryl moiety of the aryl group or arylC₁₋₄ alkyl group may be optionally substituted with a halogen atom, a C₁₋₆ alkyl group, a C₁₋₆ alkoxy group, a C₁₋₆ alkoxy carbonyl group or an amino group; R^{2''} represents hydrogen; R^{3''} represents hydrogen or C₁₋₆ alkyl; X'' represents methylene; a'' and b'' both represent 1; R^{4''} and R^{5''} both represent hydrogen; and wherein the moiety -Z''-R^{6''} represents a C₁₋₆ alkyl group, an arylC₁₋₄ alkyl group (particularly wherein aryl represents phenyl or naphthyl), a heteroarylC₁₋₄ alkyl group (particularly wherein heteroaryl represents 2-pyridyl, 3-pyridyl, 4-pyridyl or 1H-indol-3-yl), an aryloxyC₂₋₅ alkyl group or a pyrrolidinylcarbonylC₁₋₄ alkyl group in which the aryl moiety of the said groups may be optionally substituted with a halogen atom, a C₁₋₆ alkyl group, a C₁₋₆ alkoxy group, a C₁₋₆ alkoxy carbonyl group or an amino group.
- 20
- 25

Compounds of formula (I) possess a chiral carbon atom at the position marked '*' and may therefore exist as enantiomers.



5


PCT/GB01/04530 also discloses a process for the preparation of compounds of formula (I) wherein enantiomers thereof may be prepared by a combination of an achiral synthesis with a resolution step. Examples of such a resolution step are preparative chiral high performance liquid chromatography (preparative chiral HPLC) and the fractional crystallisation of diastereoisomeric salts. In particular, it is disclosed in PCT/GB01/04530 that an enantiomer of a compound of formula (I) may be prepared by the resolution of a racemic modification of a compound of formula (III)



15

wherein;


R^3 , a , b , R^4 , R^5 , Z , and R^6 are as defined in formula (I) above;
by fractional crystallisation of a diastereoisomeric salt thereof, followed by reaction of the resolved enantiomer of the compound of formula (III) with a compound of formula (X) to give a compound of formula (IV)



wherein;

L^2 and L^4 are leaving groups, and R^3 , a , b , R^4 , R^5 , Z , and R^6 are as defined in formula (I) above;

- 5 followed by reaction of a compound of formula (IV) with a compound of formula (V)




- 10 wherein;

R^1 and R^2 are as defined in formula (I) above;
to give a compound of formula (I).

Alternative processes for preparing an enantiomer of certain compounds of formula (I), being of formula (IA)


- 15



wherein;


- 20 R^1 , R^2 , b , Z , and R^6 are as defined for formula (I), and;
 k is 1 or 2;

and salts and solvates thereof has now been discovered, with the provisos that;
the compound of formula (IA) is not a compound of formula (I)^a:



wherein R^2 represents hydrogen or lower alkyl (specifically C_{1-4} alkyl); R^3 represents hydrogen; X' represents methylene or ethylene; a' and b' both represent 1; R^4 and R^5 both represent hydrogen; and wherein the moiety $-Z'-R^6$ represents halobenzyl, and;


5 the compound of formula (IA) is not a compound of formula (I)^b:



wherein $R^{1''}$ represents a hydrogen atom, a C_{1-6} alkyl group, a C_{3-6} cycloalkyl group, a C_{3-6} cycloalkyl C_{1-4} alkyl group, an aryl group or an aryl C_{1-4} alkyl group (particularly wherein aryl represents phenyl or naphthyl) in which the aryl moiety of the aryl group or aryl C_{1-4} alkyl group may be optionally substituted with a halogen atom, a C_{1-6} alkyl group, a C_{1-6} alkoxy group, a C_{1-6} alkoxy carbonyl group or an amino group; $R^{2''}$ represents a hydrogen atom or a C_{1-6} alkyl; $R^{3''}$ represents hydrogen or C_{1-6} alkyl; X'' represents methylene; a'' and b'' both represent 1; $R^{4''}$ and $R^{5''}$ both represent hydrogen; and wherein the moiety $-Z''-R^{6''}$ represents a C_{1-6} alkyl group, an aryl C_{1-4} alkyl group (particularly wherein aryl represents phenyl or naphthyl), a heteroaryl C_{1-4} alkyl group (particularly wherein heteroaryl represents 2-pyridyl, 3-pyridyl, 4-pyridyl or 1H-indol-3-yl), an aryloxy C_{2-5} alkyl group or a pyrrolidinylcarbonyl C_{1-4} alkyl group in which the aryl moiety of the said groups may be optionally substituted with a halogen atom, a C_{1-6} alkyl group, a C_{1-6} alkoxy group, a C_{1-6} alkoxy carbonyl group or an amino group.

These processes involve the chiral synthesis of certain compounds of formula (III).

25 Accordingly, in a first aspect, there is provided a process for the preparation of a compound of formula (IIIA)




or a salt thereof;

wherein;

b, Z, and R⁶ are as defined for formula (I) above, and;


k is 1 or 2;

5 which process comprises the reaction of a compound of formula (XX)



wherein;

10 b, Z, and R⁶ are as defined for formula (I);
with an enantiomer of a compound of formula (XXI)



15 wherein;

A is a ~~protection~~ amino group and k is 1 or 2;

followed by deprotection of the amino group to give a compound of formula (IIIA).

Suitable protecting groups for amines include phthalimido.


20 The compound of formula (IIIA) is typically prepared from the compounds of formulae (XX) and an enantiomer of a compound (XXI) under the Mitsunobu conditions as follows:

Typically, a mixture of the compound of formula (XX) and an enantiomer of a compound of formula (XXI) in a suitable solvent, such as tetrahydrofuran or toluene, is stirred, suitably for 8-36 hours at a suitable temperature, suitably the reflux temperature of the mixture, under an inert atmosphere, suitably an atmosphere of nitrogen. Further tetrahydrofuran is then added and the mixture cooled, suitably to 0-5°C. A phosphine, suitably triphenyl phosphine, is added and the mixture stirred until all the solid is dissolved. An *azodicarboxylate*, suitably diisopropylazodicarboxylate, is then added over a period of time, suitably 5-30 min, while maintaining the temperature at <7°C. The mixture is allowed to warm, suitably to 20-25°C. If necessary, further *phosphine and azodicarboxylate reagents* can be added. After a further period, the reaction mixture is concentrated to near dryness. A suitable alcohol, suitably propan-2-ol, is added and the concentration step repeated. This may be repeated as necessary. Further alcohol is then added and the mixture may be heated to a temperature

suitably between 65-75°C. After a suitable period, suitably 20-45 minutes, the resultant slurry is cooled to, suitably to 20-25°C, and then allowed to stand, suitably for 1.5-3 hours, after which time the product is isolated by filtration. The filter bed is washed with more alcohol and then dried *in vacuo* at 35-45°C to yield
5 the protected compound of formula (IIIA).

The protected compound of formula (IIIA) may be deprotected to yield the compound of formula (IIA) using standard conditions suitable for the removal of the particular protecting group, for example those conditions described in *P J Kocienski, Protecting Groups, (1994), Thieme*.

10 In a further aspect, the process for the preparation of the protected compound of formula (IIA) described above may also be undertaken in two stages, in which an intermediate compound of formula (IIIB);



15

wherein;

k, Z, R⁶, and b are as hereinbefore defined for formula (IIIA), and A is as hereinbefore defined for formula (XXI);
is isolated.

20 Typically, a mixture of the compound of formula (XX) and an enantiomer of a compound of formula (XXI) in a suitable solvent, such as tetrahydrofuran, C₃₋₄ alkanol, toluene, N-methylpyrrolidinone and N,N-dimethylformamide, is stirred, suitably for 8-36 hours at a suitable temperature, suitably the reflux temperature of the mixture under an inert atmosphere, suitably an atmosphere of
25 nitrogen. Further compound of formula (XX) is added as necessary and the mixture heated at a suitable temperature, suitably the reflux temperature of the mixture, under an inert atmosphere, suitably an atmosphere of nitrogen, for a suitable period of time. The reaction mixture is then cooled, suitably to 20-25°C, and the compound precipitated by means of addition of a suitable co-solvent,
30 suitably diisopropyl ether. The compound of formula (IIIB) is isolated by filtration, washed with further co-solvent and dried *in vacuo*.

A protected compound of formula (IIIA) may then be prepared from a compound of formula (IIIB) using the process described previously.


The compounds of formulae (XX) and the enantiomers of a compound of
35 formula (XXI) are known, commercially available compounds, or may be

prepared by analogy with known procedures, for examples those disclosed in standard reference texts of synthetic methodology such as *J. March, Advanced Organic Chemistry, 3rd Edition (1985), Wiley Interscience.*

The compound of formula (IIIA) is known and is disclosed in *J. Med. Chem.*, 1991, 34(2), 616-624.

The enantiomer of the compound of formula (IIIA), being the compound of formula (IIIAS) may also be isolated from a mixture of the compound of formula (IIIAS) and its antipode, typically a racemic modification, by enzymatic resolution. A mixture of the compound of formula (IIIAS) and its antipode may be prepared by using a mixture of the enantiomers of a compound of formula (XXI) in the process hereinbefore described. A racemic modification of the enantiomers of the a compound of formula (XXI) may be prepared using procedures well known in the art.

Accordingly, there is provided in a still further aspect, a process for the separation of a compound of formula (IIIAS);



wherein;

k, b, Z, and R⁶ are as hereinbefore defined for formula (IIIA);
 from its antipode, which process comprises reaction of the mixture of a compound of formula (IIIAS) and its antipode with an enzyme and a suitable enzyme donor, such as an alkyl ester of a C₄₋₈ alkanolic acid. A suitable enzyme is Lipase PS-C "Amano" II.


Typically, to a solution of a mixture of a compound of formula (IIIAS) and its antipode and a mixture of a suitable solvent, suitably *tert*-butyl methyl ether, and a suitable acyl donor, suitably ethyl octanoate, is added the enzyme, suitably Lipase PS-C "Amano" II, under an inert atmosphere, suitably an atmosphere of nitrogen. The mixture is stirred at elevated temperature, suitably 25-35°C for a suitable period of time, suitably 6-8 hours. The enzyme is removed by vacuum filtration. To the filtrate is added de-ionised water, the resultant bi-phasic solution pH adjusted to pH 4-5 and the layers separated. To the aqueous phase is added a suitable non-polar solvent, suitably dichloromethane, and the resultant bi-phasic mixture pH adjusted to pH 6-7. The layers are then separated and solvent removed in vacuo to give a compound of formula (IIIAS).

For any of the hereinbefore described reactions or processes, conventional methods of heating and cooling may be employed, for example electric heating mantles and ice/salt baths respectively.

Suitably, the absolute stereochemistry of a compound of formula (IIIA) at the position marked "*" is as shown in formula (IIIAS).


Compounds of formula (IA) may then be prepared from compounds of formula (IIIA) as follows:

The compound of formula (IIIA) is reacted with a compound of formula (XA)



wherein;

L^2 and L^4 represent leaving groups wherein L^2 and L^4 are the same or L^4 represents a leaving group which is more labile than L^2 , to form a compound of formula (IVA)



wherein;

L^2 , k, b, Z, and R^6 are as hereinbefore defined.

The compound of formula (IVA) is in turn is reacted with with a compound of formula (VA)



wherein R^1 and R^2 are as defined in formula (I) above, to give a compound of formula (IA).

Compounds of formulae (XA), and (VA) are also known, commercially available compounds, or may be prepared by analogy with known procedures, for examples those disclosed in standard reference texts of synthetic

methodology such as *J. March, Advanced Organic Chemistry, 3rd Edition (1985), Wiley Interscience.*

Suitable salts of the compounds of formula (IIIA) are those which may be useful in terms of isolation or handling of the compound of formula (IIIA) or those which may be useful in the preparation of compounds of formula (IA) and physiologically acceptable salts thereof. If appropriate, acid addition salts may be derived from inorganic or organic acids, for example tartrates, hydrochlorides, hydrobromides, sulphates, phosphates, acetates, benzoates, citrates, succinates, lactates, tartrates, fumarates, maleates, 1-hydroxy-2-naphthoates, palmoates, methanesulphonates, formates or trifluoroacetates. Salts of the compounds of formula (IIIA) may be prepared by procedures well known to those skilled in the art.

It is considered that compounds of formulae (IIIB), and (IVA) are novel. Accordingly, in an additional aspect, there is provided a compound of formula (IIIB) or a salt thereof.

There is also therefore provided a compound of formula (IVA) or a salt thereof.

Suitable salts of the compounds of the invention are those which may be useful in terms of isolation or handling of the compounds of the invention. If appropriate, acid addition salts may be derived from inorganic or organic acids, for example tartrates, hydrochlorides, hydrobromides, sulphates, phosphates, acetates, benzoates, citrates, succinates, lactates, tartrates, fumarates, maleates, 1-hydroxy-2-naphthoates, palmoates, methanesulphonates, formates or trifluoroacetates. Salts of the compounds of the invention may be prepared by procedures well known to those skilled in the art.

A compound of formula (IA) may be prepared from a compound of formula (IIIA) as follows.

Typically, a compound of formula (IIIA) in a suitable first solvent is reacted with N,N'-carbonyldiimidazole in the same solvent at reduced temperature, suitably a temperature in the range -10 - 20 °C over a suitable period of time, for example 5-60 minutes. Suitable solvents include tetrahydrofuran, dichloromethane, C₃₋₄ alkanol, isopropyl acetate, N-methylpyrrolidinone and N,N-dimethylformamide. The mixture is warmed to a suitable temperature, suitably 5-30°C and held at this temperature for a suitable period of time, for example 10-60 minutes. The compound of formula (XA) is then added, the mixture heated to a suitable elevated temperature, for example a temperature in the range 40-65°C, and stirred for a suitable period of time, for example 60-360 minutes. The reaction is then cooled to a suitable temperature, and a suitable second solvent, for example isopropyl acetate, added, followed by a aqueous solution of a suitable acidic salt, such as potassium dihydrogen

phosphate or acetic acid. The solution is clarified if necessary, the lower aqueous layer removed and the upper organic layer washed with further acidic salt solution, followed by water. The organic phase is distilled at atmospheric pressure to remove the first solvent and leave a slurry or solution of the
 5 compound of formula (IA) in the second solvent. The compound of formula (IA) may be isolated by filtration or evaporation of the solvent as appropriate.


Suitable salts of the compounds of formula (IA) include physiologically acceptable salts and salts which may not be physiologically acceptable but may be useful in the preparation of compounds of formula (IA) and physiologically
 10 acceptable salts thereof. If appropriate, acid addition salts may be derived from inorganic or organic acids, for example hydrochlorides, hydrobromides, sulphates, phosphates, acetates, benzoates, citrates, succinates, lactates, tartrates, fumarates, maleates, 1-hydroxy-2-naphthoates, palmoates, methanesulphonates, formates or trifluoroacetates. Examples of solvates include
 15 hydrates.

Salts and solvates of the compounds of formula (IA) may be prepared by procedures well known to those skilled in the art.

Suitable protecting groups in any of the above mentioned reactions are those used conventionally in the art. The methods of formation and removal of
 20 such protecting groups are those conventional methods appropriate to the molecule being protected, for example those methods discussed in standard reference texts of synthetic methodology such as *P J Kocienski, Protecting Groups, (1994), Thieme*.

Suitably, the variable R^1 of compounds of formulae (IA) and (VA)
 25 represents C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl- Y^1 -, heteroaryl- Y^1 -, aryl-(O) $_t$ -aryl- Y^1 -, aryl-(O) $_t$ -heteroaryl- Y^1 -, heteroaryl-(O) $_t$ -aryl- Y^1 -, heteroaryl-(O) $_t$ -heteroaryl- Y^1 -, aryl-SO $_2$ - Y^1 -, C_{1-6} alkyl-G- Y^1 -, J^1 -SO $_2$ - Y^1 -, R^{17} O(CO)- C_{2-6} alkenyl- Y^1 -, C_{2-6} alkynyl- Y^1 -, C_{2-6} alkenyl- Y^1 -, aryl-O- Y^1 -, heteroaryl-O- Y^1 -, C_{1-6} alkyl-SO $_2$ - Y^1 -, M- Y^1 -, J^1 - Y^1 -, J^1 -CO- Y^1 -, aryl-CO- Y^1 - or C_{3-8} cycloalkyl- Y^1 - or C_{3-8}
 30 cycloalkenyl- Y^1 -, which C_{2-6} alkynyl and C_{2-6} alkynyl- Y^1 may be optionally substituted with a -OR 17 group and which cycloalkyl or cycloalkenyl may be optionally substituted by one or more hydroxyl or C_{1-6} alkyl groups;

J^1 , J^2 and J^3 independently represent a moiety of formula (K):



35 wherein X^1 represents oxygen, NR 11 or sulphur, X^2 represents CH $_2$, oxygen, NR 12 or sulphur. m^1 represents an integer from 1 to 3 and m^2 represents an


integer from 1 to 3, provided that $m^1 + m^2$ is in the range from 3 to 5, also provided that when both X^1 and X^2 represent oxygen, NR^{11} , NR^{12} or sulphur, m^1 and m^2 must both not equal less than 2, wherein K is optionally substituted by one or more (eg. 1 or 2) $-Y^3$ -aryl, $-Y^3$ -heteroaryl, $-Y^3$ -CO-aryl, $-Y^3$ -CO-heteroaryl, $-C_{1-6}$ alkyl, $-Y^3$ -COOC $_{1-6}$ alkyl, $-Y^3$ -COC $_{1-6}$ alkyl, $-Y^3$ -W, $-Y^3$ -CO-W, $-Y^3$ -NR $^{15}R^{16}$, $-Y^3$ -CONR $^{15}R^{16}$, hydroxy, oxo, $-Y^3$ -SO $_2$ NR $^{15}R^{16}$, $-Y^3$ -SO $_2$ C $_{1-6}$ alkyl, $-Y^3$ -SO $_2$ aryl, $-Y^3$ -SO $_2$ heteroaryl, $-Y^3$ -NR $^{13}C_{1-6}$ alkyl, $-Y^3$ -NR $^{13}SO_2C_{1-6}$ alkyl, $-Y^3$ -NR $^{13}CONR^{15}R^{16}$, $-Y^3$ -NR $^{13}COOR^{14}$ or $-Y^3$ -OCONR $^{15}R^{16}$ groups, and is optionally fused to a monocyclic aryl or heteroaryl ring.

- 10 Suitably, the variable R^2 of compounds of formulae (IA) and (VA) represents hydrogen or C $_{1-6}$ alkyl.

More suitably, the variable R^1 of compounds of formulae (IA) and (VA) represents C $_{1-6}$ alkyl, C $_{2-6}$ alkenyl, C $_{2-6}$ alkynyl, aryl-Y 1 -, heteroaryl-Y 1 -, aryl-(O) $_t$ -aryl-Y 1 -, aryl-(O) $_t$ -heteroaryl-Y 1 -, heteroaryl-(O) $_t$ -aryl-Y 1 -, heteroaryl-(O) $_t$ -

- 15 heteroaryl-Y 1 -, C $_{2-6}$ alkenyl-Y 1 -, aryl-O-Y 1 -, heteroaryl-O-Y 1 -, C $_{1-6}$ alkyl-SO $_2$ -Y 1 -, M-Y 1 -, -Y 1 -J 1 , -Y 1 -CO-J 1 or C $_{3-8}$ cycloalkyl-Y 1 - or C $_{3-8}$ cycloalkenyl-Y 1 -, which cycloalkyl or cycloalkenyl may be optionally substituted by one or more hydroxyl or C $_{1-6}$ alkyl groups;

J 1 , J 2 and J 3 independently represent a moiety of formula (K):



20

wherein X^1 represents oxygen, nitrogen, NR^{11} or sulphur, X^2 represents CH $_2$, oxygen, nitrogen, NR^{12} or sulphur, m^1 represents an integer from 1 to 3, m^2 represents an integer from 1 to 3, provided that $m^1 + m^2$ is in the range from 3 to 5, also provided that when X^2 represents oxygen, nitrogen, NR^{12} or sulphur,

- 25 m^1 and m^2 must both not equal less than 2, wherein K is optionally substituted by one or more (eg. 1 or 2) $-Y^3$ -aryl, $-Y^3$ -heteroaryl, $-Y^3$ -CO-aryl, $-Y^3$ -CO-heteroaryl, $-C_{1-6}$ alkyl, $-Y^3$ -COOC $_{1-6}$ alkyl, $-Y^3$ -COC $_{1-6}$ alkyl, $-Y^3$ -W, $-Y^3$ -CO-W, $-Y^3$ -NR $^{15}R^{16}$, $-Y^3$ -CONR $^{15}R^{16}$, hydroxy, oxo, $-Y^3$ -SO $_2$ NR $^{15}R^{16}$, $-Y^3$ -SO $_2$ C $_{1-6}$ alkyl, $-Y^3$ -SO $_2$ aryl, $-Y^3$ -SO $_2$ heteroaryl, $-Y^3$ -NR $^{13}C_{1-6}$ alkyl, $-Y^3$ -NR $^{13}SO_2C_{1-6}$ alkyl, $-Y^3$ -NR $^{13}CONR^{15}R^{16}$, $-Y^3$ -NR $^{13}COOR^{14}$ or $-Y^3$ -OCONR $^{15}R^{16}$ groups, and is optionally fused to a monocyclic aryl or heteroaryl ring.

More suitably, the variable R^2 of compounds of formulae (IA) and (VA) represents hydrogen or C $_{1-6}$ alkyl.

- 35 Preferred values of Z for compounds of formulae (IIIA), (XX), and (IA) are those wherein Z represents a bond, CO, CR $^{10}R^7$ (CH $_2$) $_n$, CHR 7 (CH $_2$) $_n$ O, CHR 7 (CH $_2$) $_n$ S, CHR 7 (CH $_2$) $_n$ OCO, or CHR 7 (CH $_2$) $_n$ CO.

References to 'aryl' include references to monocyclic carbocyclic aromatic rings (eg. phenyl) and bicyclic carbocyclic aromatic rings (e.g. naphthyl) and references to 'heteroaryl' include references to mono- and bicyclic heterocyclic aromatic rings containing 1-3 hetero atoms selected from nitrogen, oxygen and sulphur. References to 'heteroaryl' may also be extended to include references to mono- and bicyclic heterocyclic aromatic rings containing 4 hetero atoms selected from nitrogen, oxygen and sulphur. Examples of monocyclic heterocyclic aromatic rings include e.g. pyridinyl, pyrimidinyl, thiophenyl, furanyl, pyrrolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, thiadiazolyl. Further examples of monocyclic heterocyclic aromatic rings include pyrazinyl, tetrazolyl or imidazolyl. Examples of bicyclic heterocyclic aromatic rings include eg. quinolinyl or indolyl. Further examples of bicyclic heterocyclic aromatic rings include benzimidazolyl. Yet further examples of bicyclic heterocyclic aromatic rings include dihydrobenzofuranyl and pyrrolopyridinyl. Carbocyclic and heterocyclic aromatic rings may be optionally substituted, e.g. by one or more C_{1-6} alkyl, C_{2-6} alkenyl, halogen, C_{1-6} alkoxy, cyano, hydroxy, nitro, amino, W, $-N(CH_3)_2$, $-NHCOC_{1-6}$ alkyl, $-OCF_3$, $-CF_3$, $-COOC_{1-6}$ alkyl, $-OCHF_2$, $-SCF_3$, $-CONR^{19}R^{20}$, $-SO_2NR^{19}R^{20}$ (wherein R^{19} and R^{20} independently represent hydrogen, C_{1-6} alkyl or C_{3-8} cycloalkyl), $-NHSO_2CH_3$, $-SO_2CH_3$ or $-SCH_3$. Yet further substituents of carbocyclic and heterocyclic aromatic rings may be $-CH_2N(CH_3)_2$ or one or more $-SH$ groups, wherein it will be appreciated that said group may tautomerise to form an $=S$ group.

Examples of group M include tetrahydronaphthalenyl.

Examples of group W include piperidinyl, pyrrolidinyl, morpholinyl and piperazinyl which may be optionally substituted with one or more C_{1-6} alkyl, halogen, or hydroxy groups.

Examples of group J^1 include $N-(COOCH_2CH_3)$ -piperidin-4-yl, $N-(CH_3)$ -piperidin-4-yl, $N-(COCH_3)$ -piperidin-4-yl, pyrrolidin-1-yl, tetrahydropyran-4-yl or N-morpholinyl. Further examples of group J^1 include $N-(cyclopropylcarbonyl)$ -piperidin-4-yl, $N-(methylsulphonyl)$ -piperidin-4-yl, thiopyranyl and tetrahydrothienyl.

Examples of group J^2 include (4-phenyl)-piperidin-1-yl, (4- $COOCH_2CH_3$)-piperazin-1-yl, (2-(3-hydroxy-pyrrolidin-1-yl-methyl))-piperidin-1-yl, N-morpholinyl, (4- $N(CH_3)_2$)-piperidin-1-yl, (4-(3-fluorophenyl))-piperazin-1-yl, (4-(4-fluorophenyl))-piperazin-1-yl, (4-pyrimidinyl)-piperazin-1-yl, (4- CH_3)-piperazin-1-yl, (4- $CONH_2$)-piperidin-1-yl, (3,3-dimethyl)-piperidin-1-yl, (4- $COCH_3$)-piperazin-1-yl, (4-(1-pyrrolidinyl-carbonylmethyl))-piperazin-1-yl, (4-hydroxy)-piperidin-1-yl, (4-methyl)-piperidin-1-yl, (4-(2-furanyl-carbonyl))-piperazin-1-yl, (4-benzyl)-

piperazin-1-yl or (3-CH₃SO₂CH₂-)-morpholin-1-yl. Further examples of group J² include thiomorpholinyl, pyrrolidinyl and benzazepinyl.

Examples of group J³ include indolinyl, which may be optionally substituted.

- 5 References to alkyl include references to both straight chain and branched chain aliphatic isomers of the corresponding alkyl. It will be appreciated that references to alkylene and alkoxy shall be interpreted similarly. References to C₃₋₈ cycloalkyl include references to all alicyclic (including branched) isomers of the corresponding alkyl.
- 10 Preferably, R¹ represents C₁₋₆ alkyl (particularly propyl), C₂₋₆ alkenyl (particularly wherein said C₂₋₆ alkenyl is substituted by one or more -COOR¹⁷ groups, eg. -HC=CH-COOH), C₂₋₆ alkynyl, aryl-Y¹-, heteroaryl-Y¹- (particularly wherein heteroaryl represents thiazolyl, indolyl, furanyl, dihydrobenzofuran, oxoimidazolyl, isoxazolyl, thienyl, thioxodihydroimidazolyl, tetrazolyl, pyrazinyl, pyrrolopyridinyl), aryl-(O)_t-aryl-Y¹-, aryl-(O)_t-heteroaryl-Y¹- (particularly wherein aryl represents phenyl and heteroaryl represents thiadiazolyl, pyrazolyl or isoxazolyl), heteroaryl-(O)_t-aryl-Y¹-, heteroaryl-(O)_t-heteroaryl-Y¹-, C₂₋₆ alkenyl-Y¹-, aryl-O-Y¹- (particularly wherein aryl represents phenyl), heteroaryl-O-Y¹-, C₁₋₆ alkyl-SO₂-Y¹- (particularly wherein C₁₋₆ alkyl represents ethyl, propyl, -CH(CH₃)₂ or -C(CH₃)₃), M-Y¹-, J¹-Y¹-, J¹-CO-Y¹-, aryl-SO₂-Y¹-, C₁₋₆ alkyl-G-Y¹- (particularly wherein C₁₋₆ alkyl represents methyl and G represents -NR¹⁸CO-, -CONR¹⁸-, -NR¹⁸SO₂- or -SO₂NR¹⁸-), heteroaryl-G-aryl-Y¹- (particularly wherein aryl represents phenyl and heteroaryl represents thiazolyl and G represents -NR¹⁸SO₂-), J¹-SO₂-Y¹- (particularly wherein J¹ represents 1-pyrrolidinyl),
- 20 R¹⁷O(CO)-C₂₋₆ alkenyl-Y¹-, R¹⁷NHCO-Y¹- (particularly wherein R¹⁷ represents hydrogen), C₂₋₆ alkynyl-Y¹- (particularly -C≡CH or wherein said C₂₋₆ alkynyl is substituted with a -OR¹⁷ group, eg. HOCH₂-C≡C-), aryl-CO-Y¹- (particularly wherein aryl represents phenyl), C₃₋₈ cycloalkyl-Y¹- or C₃₋₈ cycloalkenyl-Y¹-, which cycloalkyl or cycloalkenyl may be optionally substituted by one or more hydroxyl or C₁₋₆ alkyl groups and which C₂₋₆ alkynyl-Y¹- may be optionally substituted with a -OR¹⁷ group.
- 30 More preferred R¹ groups include C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl-Y¹-, heteroaryl-Y¹-, aryl-(O)_t-aryl-Y¹-, aryl-(O)_t-heteroaryl-Y¹-, heteroaryl-(O)_t-aryl-Y¹-, heteroaryl-(O)_t-heteroaryl-Y¹-, C₂₋₆ alkenyl-Y¹-, aryl-O-Y¹-, heteroaryl-O-Y¹-,
- 35 C₁₋₆ alkyl-SO₂-Y¹-, M-Y¹-, J¹-Y¹-, J¹-CO-Y¹- or C₃₋₈ cycloalkyl-Y¹- or C₃₋₈ cycloalkenyl-Y¹-, which cycloalkyl or cycloalkenyl may be optionally substituted by one or more hydroxyl or C₁₋₆ alkyl groups.

- Yet more preferably, R¹ represents aryl-Y¹-, heteroaryl-Y¹-, aryl-(O)_t-aryl-Y¹-, C₃₋₈ cycloalkyl-Y¹-, C₂₋₆ alkenyl-Y¹- or C₁₋₆ alkyl-SO₂-Y¹- especially wherein
- 40 aryl represents phenyl or naphthyl optionally substituted by one or more C₁₋₆

- alkyl (especially methyl), halogen (especially chlorine, fluorine and bromine), $\text{CH}_3\text{O}-$, $\text{CH}_3\text{S}-$, $\text{F}_2\text{CHO}-$, $\text{CH}_3\text{OC}(\text{O})-$, $-\text{CN}$, $-\text{CF}_3$, $\text{CF}_3-\text{S}-$, $\text{CF}_3-\text{O}-$, or $(\text{CH}_3)_2\text{N}-$, groups, and wherein heteroaryl represents pyridinyl optionally substituted by one or more halogen atoms (especially chlorine) and wherein cycloalkyl represents cyclohexyl. Further preferred substituents of phenyl include $-\text{NHCOCH}_3$ and $-\text{CONH}_2$. Yet further preferred substituents of phenyl include $-\text{SO}_2\text{NH}_2$, $-\text{CONHCH}_3$, $-\text{OCH}(\text{CH}_3)_2$, $-\text{OC}(\text{CH}_3)_3$, $-\text{COOH}$, $-\text{CON}(\text{CH}_3)_2$, $-\text{SO}_2\text{CH}_3$, $-\text{CONHCH}_2\text{CH}_3$, $-\text{CONHcyclopropyl}$ and $-\text{SO}_2\text{NHcyclopropyl}$. Also preferably, R^1 represents C_{2-6} alkynyl- Y^1 . A series of particularly preferred compounds are those wherein R^1 represents aryl- Y^1 wherein aryl represents phenyl optionally substituted by one or more C_{1-6} alkyl (especially methyl), halogen (especially chlorine, fluorine and bromine), $\text{CH}_3\text{O}-$, $\text{CH}_3\text{S}-$, $\text{F}_2\text{CHO}-$, $\text{CH}_3\text{OC}(\text{O})-$, $-\text{CN}$ or $-\text{CF}_3$ groups. Further most preferred substituents of phenyl include $-\text{NHCOCH}_3$ and $-\text{CONH}_2$. A yet further most preferred substituent of phenyl includes SO_2NH_2 . Most preferably, R^1 will also represent C_{2-6} alkenyl- Y^1 (particularly $\text{CH}_2=\text{CH}-\text{Y}^1$), C_{3-8} cycloalkyl- Y^1 (particularly cyclohexyl- Y^1) and C_{1-6} alkyl- SO_2Y^1 (particularly $\text{CH}_3\text{SO}_2-\text{Y}^1$). Also most preferably, R^1 represents C_{2-6} alkynyl- Y^1 (particularly $\text{HC}\equiv\text{C}-\text{Y}^1$).

- Especially preferred R^1 groups are aryl- Y^1 and heteroaryl- Y^1 , most especially wherein aryl represents phenyl and heteroaryl represents a 6 membered monocyclic heterocyclic aromatic ring (most particularly tetrazolyl) each of which may be optionally substituted as indicated above.

- Preferred substituents of heteroaryl include $-\text{CH}_3$, $-\text{CONH}_2$, $-\text{CH}_2\text{N}(\text{CH}_3)_2$, halogen (particularly chlorine), $-\text{OCH}_3$, $-\text{COOCH}_3$ and $-\text{NH}_2$. Most especially preferred compounds are those wherein R^1 represents phenyl- Y^1 which phenyl is substituted with a $-\text{CONH}_2$ or $-\text{CONHCH}_3$ group and tetrazolyl- Y^1 which tetrazolyl is substituted with a methyl group.

- Preferably, Y^1 represents a bond or C_{1-6} alkylene, more preferably a bond, methylene or ethylene, propylene, $-\text{C}(\text{CH}_3)_2-$ or $-\text{CH}(\text{CH}_3)-$, particularly a bond, methylene or ethylene, most preferably a bond or methylene, especially methylene.

Preferably, Y^2 represents a bond.

Preferably, Y^3 represents a bond.

- Preferably, R^2 represents hydrogen, methyl or hydroxypropyl, more preferably hydrogen or methyl, especially hydrogen.

Also preferably, R^1 and R^2 together with the nitrogen atom to which they are attached form a group of formula J^2 wherein said nitrogen atom substitutes for either X^1 or X^2 .

- Preferably, R^4 and R^5 independently represent hydrogen or methyl. Most preferably, R^4 and R^5 represent hydrogen.

Preferably, Z represents a bond, CO, $\text{CR}^{10}\text{R}^7(\text{CH}_2)_n$, $\text{CHR}^7(\text{CH}_2)_n\text{O}$, $\text{CHR}^7(\text{CH}_2)_n\text{S}$, $\text{CHR}^7(\text{CH}_2)_n\text{OCO}$ or $\text{CHR}^7(\text{CH}_2)_n\text{CO}$.

More preferably, Z represents CO, $\text{CHR}^7(\text{CH}_2)_n$, $\text{CHR}^7(\text{CH}_2)_n\text{O}$, $\text{CHR}^7(\text{CH}_2)_n\text{S}$, $\text{CHR}^7(\text{CH}_2)_n\text{OCO}$ or $\text{CHR}^7(\text{CH}_2)_n\text{CO}$, especially CH_2CO , $(\text{CH}_2)_2$,
 5 $(\text{CH}_2)_2\text{S}$, $(\text{CH}_2)_2\text{O}$, $(\text{CH}_2)_2\text{OCO}$, $(\text{CH}_2)_3\text{CO}$, CO, CHR^7 , particularly CH_2 , CHCH_3 or CH_2CO , most particularly CH_2 or CH_2CO , especially CH_2 .

Preferably, R^6 represents C_{1-6} alkyl, C_{2-6} alkenyl, CN, aryl, heteroaryl or a group of formula $-\text{Y}^2-\text{J}^3$, more preferably R^6 represents phenyl (optionally substituted with one or more halogen, phenyl or C_{2-6} alkenyl groups), naphthyl,
 10 C_{1-6} alkyl, C_{2-6} alkenyl, CN or a 5 membered aromatic heterocyclic ring containing 1 to 3 heteroatoms selected from O, N or S optionally substituted by halogen or C_{1-6} alkyl. Especially, R^6 represents phenyl (optionally substituted with one or more halogen (especially chlorine, fluorine or iodine), phenyl or 3-
 15 $\text{CH}=\text{CH}_2$ groups), naphthyl, indolyl, methyl, $-\text{CH}=\text{CH}_2$, $-\text{CN}$ or thiophenyl optionally substituted by halogen (especially chlorine). Most preferred R^6 represents indolyl (especially indolin-1-yl) or else represents phenyl substituted by one or more halogen (eg. chlorine or fluorine) groups, particularly dichlorophenyl, 3-chlorophenyl, 5-chlorothiophenyl, 4-fluorophenyl and 3,4-difluorophenyl, most particularly dichlorophenyl, especially 3,4-dichlorophenyl.

20 Preferably, R^7 represents hydrogen, methyl, COOC_{1-6} alkyl or CONR^8R^9 , more preferably hydrogen, COOC_{1-6} alkyl or CONR^8R^9 most preferably hydrogen, COOEt or CONR^8R^9 , especially hydrogen.

Preferably, R^8 and R^9 represent hydrogen.

Preferably, R^{10} represents hydrogen.

25 Preferably, R^{11} and R^{12} independently represent hydrogen or methyl.

Preferably, R^{13} and R^{14} independently represent hydrogen or methyl.

Preferably, R^{15} and R^{16} independently represent hydrogen or methyl or R^{15} and R^{16} together with the nitrogen atom to which they are attached may form a morpholine, piperidine or pyrrolidine ring, especially hydrogen or methyl.

30 Preferably, R^{17} represents hydrogen.

Preferably, R^{18} represents hydrogen.

Preferably, R^{19} and R^{20} independently represent hydrogen, C_{1-6} alkyl or C_{3-8} cycloalkyl, especially hydrogen, cyclopropyl or methyl. Particularly, R^{19} and R^{20} represent hydrogen.

35 Preferably, R^c represents hydrogen or methyl, particularly hydrogen.

Preferably, R^d represents hydrogen or methyl, particularly hydrogen.

Preferably, b represents 1.

Preferably, n represents 0, 1 or 2.

Preferably, p + q equals an integer from 0 to 2, more preferably, p and q
 40 independently represent 0 or 1 such that p + q equals an integer from 0 to 1.

Preferably, X² represents CH₂, oxygen, nitrogen or NR¹².

Preferably, J¹ represents piperidinyl (particularly piperidin-4-yl) or tetrahydropyranyl (particularly tetrahydropyran-4-yl) optionally substituted by one or two -COOCH₂CH₃, -COOtBu, -CH₃, -COCH₃, -SO₂N(CH₃)₂, -SO₂CH₃, -COPhenyl or 3, 5-dimethylisoxazol-4-ylsulphonyl groups. Also preferably, J¹ represents morpholinyl, thiopyranyl or tetrahydrothienyl which may be optionally substituted as above (particularly dioxidotetrahydrothienyl).

Preferably, J² represents piperidinyl (particularly piperidin-1-yl), morpholinyl (particularly N-morpholinyl) or piperazinyl (particularly piperazin-1-yl) optionally substituted by one or two phenyl, -COOCH₂CH₂-N(CH₃)₂, fluorophenyl, -CH₃, -CONH₂, -COCH₃, -CH₂CO-(N-pyrrolidinyl), -alkoxy, -CO-(2-furan), benzyl or -CH₂SO₂CH₃. Preferably, J² also represents thiomorpholinyl, pyrrolidinyl or benzazepinyl optionally substituted in a similar manner.

25 Preferably, J³ represents indoliny, particularly indolin-1-yl.

In a most preferred aspect, the variables R¹ and R² of compounds of formulae (IA) and (VA) represent 4-amidobenzyl and hydrogen respectively; the variables b, Z, and R⁶ for the compounds of formulae (IIIA), (XX), (IVA), and (IA) represent 1, -CH₂-, and 3,4-dichlorophenyl respectively; and the variable k for the compounds of formulae (IIIA), (XXI), and (IA) represents 1.

Suitable salts of the compounds of formula (IA) include physiologically acceptable salts and salts which may not be physiologically acceptable but may be useful in the preparation of compounds of formula (I) and physiologically acceptable salts thereof. If appropriate, acid addition salts may be derived from inorganic or organic acids, for example hydrochlorides, hydrobromides, sulphates, phosphates, acetates, benzoates, citrates, succinates, lactates, tartrates, fumarates, maleates, 1-hydroxy-2-naphthoates, palmoates, methanesulphonates, formates or trifluoroacetates. Examples of solvates include hydrates. Salts and solvates of the compounds of formula (IA) may be prepared by procedures well known to those skilled in the art.

Throughout the specification and the claims which follow, unless the context requires otherwise, the word 'comprise', and variations such as 'comprises' and 'comprising', will be understood to imply the inclusion of a stated integer or step or group of integers but not to the exclusion of any other integer or step or group of integers or steps.

The following Examples illustrate the invention but do not limit it in any way.

General experimental details

10

NMR

Nuclear magnetic resonance (NMR) spectra were acquired using a Bruker DPX250 or DPX400 instrument.

15 IR

Infra-red spectra were acquired using a Nicolet Avatar 360 instrument using a Germanium ATR probe

LC/MS System

- 20 The following Liquid Chromatography-Mass Spectroscopy (LCMS) system was used: 3mm ABZ+PLUS (3.3cm x 4.6mm internal diameter) column, eluting with solvents: A – 0.1% formic acid + 0.077% w/v ammonium acetate in water; and B – 95:5 acetonitrile:water + 0.05%v/v formic acid, at a flow rate of 3ml per minute. The following gradient profile was used: 100% A for 0.7min; A + B mixtures, gradient profile 0 – 100% B over 3.5min; hold at 100%B for 1.1min; return to 100% A over 0.2min.

Analytical HPLC column, conditions and eluent

- Reverse-phase high performance liquid chromatography was carried out using a Luna 3mm C18(2) (50 x 2.0mm i.d.) column eluting with solvents: A – 100% water, 0.05% TFA; and B – 100% acetonitrile, 0.05%TFA, at a flow rate of 2ml per minute, and at 60°C. The following gradient profile was used:0-95% B over 2.00min, return to 0% B over 0.01min.

35 Examples

Example 1: Preparation of [(2S)-4-(3,4-dichlorobenzyl)morpholin-2-yl]methylamine – enzyme method

- To a solution of [4-(3,4-dichlorobenzyl)morpholin-2-yl]methylamine (3g) and ethyl octanoate (5.8ml) in *tert*-butyl methyl ether (30ml) was added enzyme Lipase PS-C "Amano" II (3g), under an atmosphere of nitrogen. The mixture was stirred

at 200rpm and heated to 30°C. The mixture was stirred at 30°C for a further 6.5h. The enzyme was removed by vacuum filtration. To the filtrate was added de-ionised water (15ml). The resultant bi-phasic solution was pH adjusted to pH 5.5 and the layers were separated. To the aqueous phase was added DCM (15ml) and the resultant bi-phasic mixture was pH adjusted to pH 6.5. The layers were separated and solvent was evaporated *in vacuo* to give the title compound as a yellow oil (1.0g, 98.9%a/a, 94.8%ee)
LC/MS (System A) R_t 1.77 min, Mass Spectrum *m/z* 275 [MH⁺].

10 Example 2 - Preparation of 2-[(2*R*)-4-(3,4-dichlorobenzyl)morpholin-2-yl]methyl]-1*H*-isoindole-1,3(2*H*)-dione

A mixture of 2-[(3,4-dichlorobenzyl)amino]ethanol (2.038 g) and (S)-2-(oxiran-2-ylmethyl)-1*H*-isoindole-1,3(2*H*)-dione (N-(2,3-epoxypropyl)-phthalimide) (2.032g) in tetrahydrofuran (3.3ml) was stirred and heated at reflux under nitrogen. After 21.5h more tetrahydrofuran (12.5ml) was added and the mixture was cooled to 3°. Triphenyl phosphine (2.793g) was added and the mixture was stirred until all the solid had dissolved. Diisopropylazodicarboxylate (2.1ml) was then added over 12min maintaining the temperature at <7°. After 2.25h the mixture was allowed to warm to 22°. After 5.3h more triphenylphosphine (121mg) and diisopropylazodicarboxylate (0.09ml) were added. After 22.5h the reaction mixture was concentrated to near dryness. Propan-2-ol (12ml) was added and the concentration repeated, this was repeated once more. More propan-2-ol (12ml) was added and the mixture was heated to 70°. After 0.5h the slurry was cooled to 22° and then after a further 2h the product was collected. The bed was washed with propan-2-ol (2x4ml) and then dried *in vacuo* at 40° to give the title compound (2.622g).

NMR (DMSO *d*-6): 1.93δ (1H) d of d, J=11.0Hz, 8.8Hz; 2.10δ (1H) d of t, J=3.5Hz, 11.3Hz; 2.52δ (1H) broad d, J=11.3Hz; 2.77δ (1H) broad d, J=11.3Hz; 3.3 – 3.8δ (7H) m; 7.31δ (1H) d of d, J=8.2Hz, 1.9Hz; 7.55δ (1H) d, J=1.9Hz; 7.68δ (1H) d, J=8.2Hz; 7.86δ (4H) m.

Preparation of [(2*S*)-4-(3,4-dichlorobenzyl)morpholin-2-yl]methylamine

A slurry of 2-[(2*R*)-4-(3,4-dichlorobenzyl)morpholin-2-yl]methyl]-1*H*-isoindole-1,3(2*H*)-dione (1.00g) in water(8.5ml) was heated to 75° and then treated dropwise with concentrated sulphuric acid (2.5ml). The mixture was then heated at reflux. After 23h the reaction mixture was cooled to 22° and then treated with dichloromethane (6ml). 880 Ammonia solution (7ml) was then added dropwise with cooling. More dichloromethane (10ml) was added. The aqueous phase was separated and extracted with more dichloromethane (10ml). The combined organic phase was washed with water (5ml) and then evaporated to dryness.

The residue was reevaporated from DCM to give the title compound as an oil (662mg).

LC/MS (System A) R_t 1.77 min, Mass Spectrum m/z 275 [MH^+].

5 Description 1: Preparation of 2-[(2*R*)-3-[(3,4-dichlorobenzyl)(2-hydroxyethyl)amino]-2-hydroxypropyl]-1*H*-isoindole-1,3(2*H*)-dione

- To a solution of 2-[(3,4-dichlorobenzyl)amino]ethanol (2.8g) in tetrahydrofuran (6.2 ml) is added (S)-2-(oxiran-2-ylmethyl)-1*H*-isoindole-1,3(2*H*)-dione (N-(2,3-epoxypropyl)-phthalimide) (3.1g) with stirring, under a nitrogen atmosphere. The mixture was heated to 90 °C over 1 h, then held at this temperature for 18 h. Further 2-[(3,4-dichlorobenzyl)amino]ethanol (0.14g) is added, and the reaction mixture heated to 90 °C for a further 5h. The reaction mixture is cooled to 22 °C, and diisopropyl ether (21ml) added, and the product isolated by vacuum filtration. The filter cake is washed with diisopropyl ether (1 vol) and dried in vacuo at 40 ° to give the title compound as a white solid.

LC/MS System B

- 3µm Phenomenex Luna (50 x 2mm i.d.) column, eluting with solvents: A – 0.05% trifluoroacetic acid in water, B – 0.05% trifluoroacetic acid in acetonitrile, at 40°C and at a flow rate of 1ml per minute. The following linear gradient was used: 0 to 95% B over 8 minutes.

LC/MS (System B) R_t 3.85min, Mass Spectrum m/z 423 [MH^+]

Description 2: Preparation of 4-([[(2*S*)-4-(3,4-dichlorobenzyl)morpholin-2-yl]methyl]amino)carbonyl]amino)methyl)benzamide benzenesulfonate hydrate

- 25 A solution of [(2*S*)-4-(3,4-dichlorobenzyl)morpholin-2-yl]methylamine (5g) in THF (10ml) is added to a slurry of N,N'-carbonyldiimidazole (3.2g) in THF (30ml) at 5-10 °C over ca. 10 min. The mixture is warmed to 15±3° and held at this temperature for ca. 15min. 4-Aminomethyl benzamide (3.0g) is then added, the mixture heated to 60±3° and stirred at this temp for 75 min.
- 30 The reaction is cooled to 22±3° and isopropyl acetate (40ml) added, followed by a solution of potassium dihydrogen phosphate (5% w/v, 40ml). The solution is filtered through celite (2g), the lower aqueous layer is removed and the upper organic layer washed with potassium dihydrogen phosphate (5% w/v, 2x40ml) then water (40ml). The organic phase is distilled at atmospheric pressure to
- 35 remove THF and leave a slurry of 4-([[(2*S*)-4-(3,4-dichlorobenzyl)morpholin-2-yl]methyl] amino)carbonyl]amino)methyl)benzamide in isopropyl acetate (ca 60ml).
- This is cooled to 50±3° and isopropanol (30ml) is added, followed by an aqueous solution of benzene sulfonic acid (32% w/v, 10ml). The mixture is cooled to
- 40 22±3° over ca 1h, seeded with authentic 4-([[(2*S*)-4-(3,4-

dichlorobenzyl)morpholin-2-yl)methyl} amino)carbonyl]amino)methyl)benzamide hydrate and aged at $22 \pm 3^\circ$ for 72 h. The contents are cooled to $0 \pm 3^\circ$ over 1 h and filtered. The filter cake is washed with a 4:1:0.1 mixture of isopropyl acetate/isopropyl alcohol/water (2.5ml) and dried in vacuo at $25 \pm 5^\circ$ to give the
5 title compound as a white solid (6.9g).

NMR (DMSO d-6): 2.81 δ (1H) broad t; 3.0 – 3.4 δ (5H) m; 3.67 δ (2H) m; 4.02 δ (1H) d of d, J=12.7Hz, 2.5Hz; 4.25 δ (1H) d, 5.9Hz; 4.37 δ (2H) m; 6.24 δ (1H) t, J=5.6Hz; 6.58 δ (1H) t, J=5.9Hz; 7.3 δ (6H) m; 7.48 δ (1H) d of d, J=8.3Hz, 2.0Hz; 7.61 δ (2H) m [benzene sulphonate]; 7.75 δ (1H) d, J=8.3Hz; 7.81 δ (1H) d, 2.0Hz;
10 7.82 δ (2H) m; 7.91 δ (1H) broad s; 9.85 (1H) broad s [NH⁺].